

First Total Synthesis of 1,3-Diacetyl- and -Dibutyroyl-2-oleoylglycerol, Previously Isolated From Natural Products

J. GUILLON, B. LELONG, J. -P. RIOULT* AND M. ROBBA

Syntheval, 5 rue Vaubénard, 14032 Caen Cedex and *Laboratoire de Botanique et de Mycologie, UFR des Sciences Pharmaceutiques, Boulevard Becquerel, 14032 Caen Cedex, France

Abstract

The first total synthesis of 1,3-diacetyl-2-oleoylglycerol and 1,3-dibutyroyl-2-oleoylglycerol is described in two steps starting from 1,3-dichloropropan-2-ol. These two glycerides were previously described by hemi-synthesis from natural products.

We describe the total synthesis of two symmetrical triglycerides, 1,3-diacetyl-2-oleoylglycerol (**1a**) and 1,3-dibutyroyl-2-oleoylglycerol (**1b**) (Figure 1), starting from commercially available 1,3-dichloropropan-2-ol (**2**). Compounds **1a** and **1b** could be potentially interesting in the prevention of cardiovascular diseases.

The 1,3-acetylated triglyceride **1a** was first isolated from a yeast, *Lipomyces starkeyi* Lodder & Kreger-van Rij (Smith et al 1995), grown in a medium containing 1,2-propanediol followed by acetolysis (Suzuki & Hasegawa 1974). In our work, the 1,3-butyroyl triglyceride **1b** and the 1,3-acetylated triglyceride **1a** were prepared in two steps by hemi-synthesis from a microorganism oil rich in docosahexaenoic acid (Bayon & Dolmazon 1996). The oil was a clear yellow-orange liquid (triglyceride) prepared from a marine dinoflagellate, *Cryptocodinium cohnii* (Seligo) Javornicky (Javornicky 1962), which is a non-photosynthetic and non-toxic marine dinoflagellate rich in triacylglycerols (Henderson et al 1988; Henderson & Mackinlay 1991). This heterotrophic marine protoctista (Margulis 1988) is principally localized on rotting *Fucus thalli*, on the littoral of the Baltic Sea, the Atlantic Ocean and the Mediterranean Sea.

Materials and Methods

Chemistry

Melting points were determined on a Kofler block and are uncorrected. IR spectra were recorded on a Genesis Series FTIR spectrometer. ¹H and ¹³C

NMR spectra were recorded on a Jeol JNM-LA 400 spectrometer (400 MHz) using CDCl₃ or d₆-DMSO as solvent. Chemical shifts refer to tetramethylsilane which was used as an internal reference. Key: s = singlet, d = doublet, t = triplet, dd = double doublet, qt = quintuplet, sx = sextuplet, m = multiplet. Mass spectra were recorded on a Jeol D 300 instrument using a direct inlet system and electron-impact ionization. Silica gel 60 (70–230 mesh) was used for column chromatography. Elemental analyses (C, H) were performed by INSA (Rouen, France), and agreed with the proposed structures within ±0.3% of the theoretical values.

2-Chloro-1-(chloromethyl)ethyl oleate (**3**)

The reaction between 1,3-dichloropropan-2-ol (**2**) (15 g, 0.116 mol) and oleoyl chloride (30.09 g, 0.10 mol) was completed by heating the mixture at 100°C for 4 h under stirring. The reaction mixture was then cooled and extracted with diethyl ether. The organic layer was washed with water, twice with aqueous sodium bicarbonate, dried over sodium sulphate, charcoalated and evaporated to dryness to give **3** as an orange oil (yield 69%); IR (KBr) ν : 1745 cm⁻¹ (CO); ¹H NMR (CDCl₃) δ : 0.87 (3H, t, J = 6.77, CH₃ oleoyl), 1.28 (22H, m, 11 CH₂ oleoyl), 1.64 (2H, m, CH=CH-CH₂), 1.96 (1H, m, CH₂-CH=CH), 2.00 (1H, m, CH₂-CH=CH), 2.37 (2H, t, J = 7.50, CH₂-CO), 3.74 (2H, d, J = 5.14, CH₂-CH-CH₂), 3.76 (2H, d, J = 5.14, CH₂-CH-CH₂), 5.18 (1H, qt, J = 5.14, CH₂-CH-CH₂), 5.35 (1H, m, CH=CH), 5.38 (1H, m, CH=CH).

Correspondence: J. Guillon, Syntheval, 5 rue Vaubénard, 14032 Caen Cedex, France




Figure 1 General structure of compounds 1a and 1b.

General procedure for preparation of silver acetate (4a) and silver butyrate (4b)

To a solution of acetic acid or butyric acid (0.17 mol) in 500 mL ethanol, were added 28% aqueous ammonia (23.7 mL, 0.17 mol) and silver nitrate (28.92 g, 0.17 mol) dissolved in twice its weight of water. After stirring for 4 h at room temperature, the precipitate thus formed was filtered off, washed with ethanol and dried at 60°C for several hours.

Silver butyrate (4b). White-grey crystals (yield 51%); mp 254°C; IR (KBr) ν : 1565 cm⁻¹ (COOAg); ¹H NMR (d₆-DMSO) δ : 0.88 (3H, t, J = 7.42, CH₃), 1.54 (2H, sx, J = 7.42, CH₂), 2.13 (2H, t, J = 7.42, CH₂-CO); ¹³C NMR (d₆-DMSO) δ : 13.91 (CH₃), 19.47 (CH₂), 38.65 (CH₂), 177.27 (COOAg).

General procedure for preparation of 1,3-diacetyl-2-oleoylglycerol (1a) and 1,3-dibutyryl-2-oleoylglycerol (1b)

Silver acetate (4a) or silver butyrate (4b) (0.0152 mol) were mixed with 2-chloro-1-(chloromethyl)ethyl oleate (3) (0.00508 mol), with the aid of diethyl ether. The ether was removed by warming on a water-bath, and the mixture heated at 170°C for 8 h with occasional stirring. When cold, the product was extracted with diethyl ether and silver chloride was filtered off. The organic layer was then washed twice with aqueous sodium bicarbonate, then with water, dried over magnesium sulphate, charcoal and evaporated to dryness. The brown oily residue was twice chromatographed on a silica-gel column, eluting with diethyl ether-hexane (20:80, v/v) for the first column and with ethyl acetate-hexane (20:80, v/v) for the second.

1,3-Diacetyl-2-oleoylglycerol (1a). Colourless oil (yield 30%); IR (KBr) ν : 1745 cm⁻¹ (CO); ¹H NMR (CDCl₃) δ : 0.81 (3H, t, J = 6.75, CH₃ oleoyl), 1.21 (22H, m, 11 CH₂ oleoyl), 1.53 (2H, m,

CH=CH-CH₂), 1.89 (1H, m, CH₂-CH=CH), 1.94 (1H, m, CH₂-CH=CH), 2.01 (6H, s, 2 CH₃CO acetyl), 2.26 (2H, t, J = 7.40, CH₂-CO oleoyl), 4.08 (2H, dd, J = 11.90 and 5.90, CH₂-CH-CH₂), 4.21 (2H, dd, J = 11.90 and 4.15, CH₂-CH-CH₂), 5.19 (1H, m, CH₂-CH-CH₂), 5.27 (1H, m, CH=CH), 5.30 (1H, m, CH=CH); ¹³C NMR (CDCl₃) δ : 14.04 (CH₃ oleoyl), 20.69 (2 CH₃-CO acetyl), 22.62 (CH₂), 24.77 (CH₂), 27.08 (CH₂), 27.15 (CH₂), 29.03 (CH₂), 29.11 (CH₂), 29.25 (CH₂), 29.29 (CH₂), 29.41 (CH₂), 29.45 (CH₂), 29.59 (CH₂), 29.68 (CH₂), 32.20 (CH₂), 34.02 (CH₂-CO oleoyl), 62.10 (2 COO-CH₂), 68.84 (COO-CH), 129.78 (CH=CH), 130.40 (CH=CH), 170.40 (2 CO acetyl), 173.02 (CO oleoyl); m/z (EI) 440 (M⁺), other major fragments; 380, 264, 227, 171, 159, 137, 117, 98. Calculated for C₂₅H₄₄O₆: C, 68.08; H, 9.98. Found: C, 68.25; H, 10.11%.

1,3-Dibutyryl-2-oleoylglycerol (1b). Pale yellow oil (yield 21%); IR (KBr) ν : 1745 cm⁻¹ (CO); ¹H NMR (CDCl₃) δ : 0.88 (3H, t, J = 6.75, CH₃ oleoyl), 0.95 (6H, t, J = 7.45, 2 CH₃ butyryl), 1.29 (22H, m, 11 CH₂ oleoyl), 1.63 (2H, m, CH=CH-CH₂), 1.67 (4H, sx, J = 7.45, CH₂ butyryl), 1.97 (1H, m, CH₂-CH=CH), 2.01 (1H, m, CH₂-CH=CH), 2.30 (4H, t, J = 7.45, 2 CH₂-CO butyryl), 2.31 (2H, t, J = 7.30, CH₂-CO oleoyl), 4.15 (2H, dd, J = 11.88 and 5.93, CH₂-CH-CH₂), 4.30 (2H, dd, J = 11.88 and 4.33, CH₂-CH-CH₂), 5.27 (1H, m, CH₂-CH-CH₂), 5.34 (1H, m, CH=CH), 5.38 (1H, m, CH=CH); ¹³C NMR (CDCl₃) δ : 13.58 (2 CH₃ butyryl), 14.08 (CH₃ oleoyl), 18.31 (2 CH₂ butyryl), 22.66 (CH₂), 24.85 (CH₂), 27.14 (CH₂), 27.19 (CH₂), 29.04 (CH₂), 29.16 (CH₂), 29.24 (CH₂), 29.26 (CH₂), 29.49 (CH₂), 29.62 (CH₂), 29.73 (CH₂), 31.87 (CH₂), 32.57 (CH₂), 34.16 (CH₂-CO oleoyl), 35.88 (2 CH₂-CO butyryl), 62.05 (2 COO-CH₂), 68.83 (COO-CH), 129.66 (CH=CH), 130.00 (CH=CH), 172.84 (CO oleoyl), 173.07 (2 CO butyryl); m/z (EI) 496 (M⁺), other major fragments; 408, 395, 337, 264, 215, 145, 98, 71. Calculated for C₂₉H₅₂O₆: C, 70.12; H, 10.55. Found: C, 70.29; H, 10.78%.

Results and Discussion

Chemistry

The reaction sequence used to prepare 1,3-diacetylated and -butyrylated triglycerides (1a and 1b)




Figure 2. Synthesis of 1,3-diacyl-2-oleoylglycerol (1a) and 1,3-dibutyryl-2-oleoylglycerol (1b). Reagents: i. (Z)- $CH_3-(CH_2)_7-CH=CH-(CH_2)_7-COCl$; ii. $CH_3-COOAg$ or $CH_3-(CH_2)_2-COOAg$.




Figure 3. Synthesis of silver acetate (4a) and silver butyrate (4b). Reagent: i. $AgNO_3$, NH_4OH , C_2H_5OH .

began from commercially available 1,3-dichloropropan-2-ol (2) and was achieved in two steps with 21 and 15% overall yield, respectively. 2-Chloro-1-(chloromethyl)ethyl oleate (3) (Humnicki 1929; Bougault 1932; Schuster 1933) was obtained in 69% yield by heating oleoyl chloride at 100°C with 1,3-dichloropropan-2-ol (2) according to Grün's method (Grün 1910). The derivative (3) was subjected to nucleophilic substitution with 3 molar equivalent, silver acetate (4a) or silver butyrate (4b). This led to symmetrical 1,3-diacyl-2-oleoylglycerol (1a) and 1,3-dibutyryl-2-oleoylglycerol (1b) (Whitby 1926; Bhati et al 1983), respectively. The silver salts of acetic or butyric acids (4a or 4b) reacted very readily with the alkyl dihalide (3) yielding triesters 1a and 1b (Figure 2).

Silver acetate (4a) and silver butyrate (4b) (Peterson et al 1952) were prepared by reaction of silver nitrate with acetic acid or butyric acid in the presence of aqueous ammonia in ethanol (Whitby 1926; Figure 3).

References

- Rayon, Y., Dolmazan, R. (1996) Two step synthesis of 1,3-acetylated, -butyrylated and -caproylated triglycerides from a microorganism oil rich in docosahexaenoic acid (DHA). *Bioorg. Med. Chem. Lett.* 6: 345-348
- Bhati, A., Hamilton, R. J., Steven, D. A., Ancja, R., Padley, F. B. (1983) Synthesis of triglycerides from 1,3-dibromopropan-2-ol. *J. Chem. Soc. Perkin Trans. 2*: 1553-1558
- Bougault, M. J. (1932) Synthèse de l' α -x distéaro- β -azélaïne. *J. Pharm. Chim.* 16: 428-431
- Grün, A. (1910) Synthese der symmetrischen Monoglyceride. *Chem. Ber.* 43: 1288-1291
- Henderson, R. J., Mackinlay, E. E. (1991) Polyunsaturated fatty acid metabolism in the marine dinoflagellate *Cryptocodinium cohnii*. *Phytochemistry* 30: 1781-1787
- Henderson, R. J., Lettley, J. W., Sargent, J. R. (1988) Lipid composition and biosynthesis in the marine dinoflagellate *Cryptocodinium cohnii*. *Phytochemistry* 27: 1679-1683
- Humnicki, V. (1929) Sur les glycérides mixtes de l'acide salicylique (Ile partie). *Bull. Soc. Chim. Fr.* 45: 279-283
- Javornicky, P. (1962) Two scarcely known genera of the class Dinophyceae: *Bernardinium* Chodai and *Cryptocodinium* Biecheler. *Preslia (Pratia)* 34: 98-113
- Margulis, L. (1988) *Protozoists & Dinoflagellata*. In: Margulis, L., Schwartz, K. V. (eds) *Five Kingdoms, An Illustrated Guide to the Phyla of Life on Earth*. 2nd edn. Freeman, W. H. & Co, New York, pp 77-83
- Peterson, S., Jones, J. L., Schmitt, J. M., Knabeschi, L. H. (1952) The solubility of silver propionate and silver butyrate in aqueous sodium nitrate. *J. Phys. Chem.* 56: 491-492
- Schuster, G. (1933) Zusammensetzung der Illipébutter. Untersuchung über α -Dilauryl- β -azélaïn und α , α -Dilaurin. *Chem. Zentralbl.* 104: 1702
- Smith, M. T., van der Walt, J. P., Batenburg-van der Vegte, W. H. (1995) *Bafjevia* gen. nov. - a new genus of the Lipomycetaceae. *Antonie van Leeuwenhoek* 67: 177-179
- Suzuki, T., Hasegawa, K. (1974) Diol lipids in the phospholipid fraction of *Lipomyces starkeyi* grown in the medium containing 1,2-propanediol. *Agr. Biol. Chem.* 38: 613-620
- Whitby, G. S. (1926) Some fatty acid derivatives. *J. Chem. Soc.* 128: 1458-1465